

Symbolic Parameter Estimation of Continuous-Time Markov Chains

Anders Malta Jakobsen
dept. of Computer Science
AAU
Aalborg, Denmark
amja23@student.aau.dk

Casper Ståhl
dept. of Computer Science
AAU
Aalborg, Denmark
cstahl20@student.aau.dk

Daniel Runge Petersen
dept. of Computer Science
AAU
Aalborg, Denmark
dpet20@student.aau.dk

Lars Emanuel Hansen
dept. of Computer Science
AAU
Aalborg, Denmark
leha20@student.aau.dk

Oliver Holmgaard
dept. of Computer Science
AAU
Aalborg, Denmark
oholmg20@student.aau.dk

Sebastian Aaholm
dept. of Computer Science
AAU
Aalborg, Denmark
saahol20@student.aau.dk

Abstract—This is a placeholder abstract. The whole template is used in semester projects at Aalborg University (AAU).

I. INTRODUCTION

This paper is about improving the runtime of jajapy - a tool for estimating parameters in parametric models.

Markov Chain (MC) - A chain of events described as a sequence of events without knowledge of prior. Hidden Markov Model (HMM) - A markov chain with emission probabilities. Markov Decision Process (MDP) - A markov chain with actions that influence the transitions. Continuous Time Markov Chain - A markov chain with traces that have dwell times as well as label emissions. Baum-Welch algorithm (BW) - Expectation-Maximization algorithm for finding the parameters of a Hidden Markov Model. Algebraic Decision Diagram (ADD) - A data structure of states and binary decisions, also called a Multi-Terminal Binary Decision Diagram (MTBDD).

II. HMM EXAMPLE

A. Setup

We have a simple HMM with, two hidden states S_1 and S_2 , two observation symbols: O_1 and O_2 and an observation sequence $O = \{O_1, O_2, O_1\}$.

The HMM parameters are:

Transition matrix A (probability of moving from one state to another):

$$A = \begin{bmatrix} 0.6 & 0.4 \\ 0.5 & 0.5 \end{bmatrix}$$

Emission matrix B (probability of emitting observation given a state):

$$B = \begin{bmatrix} 0.7 & 0.3 \\ 0.4 & 0.6 \end{bmatrix}$$

Initial state probability vector π (probability of starting in each state):

$$\pi = [0.8 \quad 0.2]$$

B. Expectation step

In the expectation step we calculate α and β .

1) *Forward step* α : We first compute the forward probabilities $\alpha_t(i)$, which represent the probability of being in state i at time t after observing the first t symbols.

a) *Initialization* at $(t = 1)$:

$$\alpha_1 = \pi \circ B_{y_1}$$

Where B_{y_1} is the first column of the emission matrix, corresponding to observation O_1

(i.e., $B_{y_1} = \begin{bmatrix} 0.7 \\ 0.4 \end{bmatrix}$) and \circ represents the Hadamard product.

So, we get:

$$\alpha_1 = \begin{bmatrix} 0.8 \\ 0.2 \end{bmatrix} \circ \begin{bmatrix} 0.7 \\ 0.4 \end{bmatrix} = \begin{bmatrix} 0.56 \\ 0.08 \end{bmatrix}$$

b) *Induction* (for $t = 2, 3, \dots, T$): For subsequent timesteps, we compute:

$$\alpha_{t+1} = B_{y_{t+1}} \circ (A^T \alpha_t)$$

Where A^T is the transpose of the transition matrix. Let's apply this to compute the forward probabilities for $t = 2$ and $t = 3$:

At $t = 2$ (observation O_2):

$$\alpha_2 = B_{y_2} \circ (A^T \alpha_1)$$

We have:

$$B(y_2) = \begin{bmatrix} 0.3 \\ 0.6 \end{bmatrix}$$

and

$$A^T = \begin{bmatrix} 0.6 & 0.5 \\ 0.4 & 0.5 \end{bmatrix}$$

We get:

$$\alpha_2 = \begin{bmatrix} 0.3 \\ 0.6 \end{bmatrix} \circ \left(\begin{bmatrix} 0.6 & 0.5 \\ 0.4 & 0.5 \end{bmatrix} \cdot \begin{bmatrix} 0.56 \\ 0.08 \end{bmatrix} \right) = \begin{bmatrix} 0.1128 \\ 0.1584 \end{bmatrix}$$

At $t = 3$ (observation O_1):

$$\alpha_2 = B_{y1} \circ (A^T \alpha_2)$$

We get:

$$\alpha_3 = \begin{bmatrix} 0.7 \\ 0.4 \end{bmatrix} \circ \left(\begin{bmatrix} 0.6 & 0.5 \\ 0.4 & 0.5 \end{bmatrix} \cdot \begin{bmatrix} 0.1 \\ 0.1584 \end{bmatrix} \right) = \begin{bmatrix} 0.102816 \\ 0.049728 \end{bmatrix}$$

2) *Backward step β* : The backward probabilities $\beta_t(i)$ represent the probability of observing the rest of the sequence starting from time $t + 1$, given that the system is in state i at time t .

Initialization (at $t = T = 3$)

$$\beta_T = \mathbf{1} = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

a) *Induction (for $t = T - 1, T - 2, \dots, 1$)*: For earlier timesteps, we compute:

$$\beta_t = A(\beta_{t+1} \circ B_{y_{t+1}})$$

At $t = 2$ (observation O_1):

$$\beta_2 = A(\beta_3 \circ B_{y1})$$

$$B_{y1} = \begin{bmatrix} 0.7 \\ 0.4 \end{bmatrix}, \quad \beta_3 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

We get:

$$\beta_2 = \begin{bmatrix} 0.6 & 0.4 \\ 0.5 & 0.5 \end{bmatrix} \cdot \left(\begin{bmatrix} 1 \\ 1 \end{bmatrix} \circ \begin{bmatrix} 0.7 \\ 0.4 \end{bmatrix} \right)$$

$$\beta_2 = \begin{bmatrix} 0.6 & 0.4 \\ 0.5 & 0.5 \end{bmatrix} \cdot \begin{bmatrix} 0.7 \\ 0.4 \end{bmatrix} = \begin{bmatrix} 0.58 \\ 0.55 \end{bmatrix}$$

At $t = 1$ (observation O_2):

$$\beta_1 = A(\beta_2 \circ B_{y2})$$

We have:

$$B_{y2} = \begin{bmatrix} 0.3 \\ 0.6 \end{bmatrix}, \quad \beta_2 = \begin{bmatrix} 0.58 \\ 0.55 \end{bmatrix}$$

$$\beta_1 = \begin{bmatrix} 0.6 & 0.4 \\ 0.5 & 0.5 \end{bmatrix} \cdot \left(\begin{bmatrix} 0.3 \\ 0.6 \end{bmatrix} \circ \begin{bmatrix} 0.58 \\ 0.55 \end{bmatrix} \right)$$

$$\beta_1 = \begin{bmatrix} 0.6 & 0.4 \\ 0.5 & 0.5 \end{bmatrix} \cdot \begin{bmatrix} 0.174 \\ 0.33 \end{bmatrix} = \begin{bmatrix} 0.2364 \\ 0.252 \end{bmatrix}$$

C. *Step 3: Compute γ and ξ*

1) *Compute γ* : We can compute γ by

$$\gamma_t = (\mathbb{1}^T \cdot \alpha_T)^{-1} \cdot (\alpha_t \circ \beta_t)$$

$$\alpha_T = \begin{bmatrix} 0.089628 \\ 0.053328 \end{bmatrix}$$

$$\mathbb{1}^T \cdot \alpha_T = 0.089628 + 0.053328 = 0.152544$$

This is the total probability of observing our sequence $O = \{O_1, O_2, O_1\}$

Now we can compute γ_t for each time stamp.

At $t=1$: We have

$$\alpha_1 = \begin{bmatrix} 0.56 \\ 0.08 \end{bmatrix}, \quad \beta_1 = \begin{bmatrix} 0.2364 \\ 0.252 \end{bmatrix}$$

We take the Hadamard product of this.

$$\alpha_1 \circ \beta_1 = \begin{bmatrix} 0.56 \cdot 0.2364 \\ 0.08 \cdot 0.252 \end{bmatrix} = \begin{bmatrix} 0.132384 \\ 0.02016 \end{bmatrix}$$

We normalize the first part and take the scalar product.

$$\gamma_1 = \frac{1}{0.152544} \cdot \begin{bmatrix} 0.132384 \\ 0.02016 \end{bmatrix} = \begin{bmatrix} 0.8678414 \\ 0.1321589 \end{bmatrix}$$

At $t = 2$:

We have:

$$\alpha_2 = \begin{bmatrix} 0.1074 \\ 0.1584 \end{bmatrix}, \quad \beta_2 = \begin{bmatrix} 0.58 \\ 0.55 \end{bmatrix}$$

The Hadamard product is:

$$\alpha_2 \circ \beta_2 = \begin{bmatrix} 0.1074 \cdot 0.58 \\ 0.1584 \cdot 0.55 \end{bmatrix} = \begin{bmatrix} 0.062292 \\ 0.08712 \end{bmatrix}$$

We normalize the first part and take the scalar product.

$$\gamma_2 = \frac{1}{0.152544} \cdot \begin{bmatrix} 0.062292 \\ 0.08712 \end{bmatrix} = \begin{bmatrix} 0.42888609 \\ 0.57111391 \end{bmatrix}$$

At $t = 3$

We have:

$$\alpha_3 = \begin{bmatrix} 0.089628 \\ 0.053328 \end{bmatrix}, \quad \beta_3 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

The Hadamard product is:

$$\alpha_3 \circ \beta_3 = \begin{bmatrix} 0.089628 \\ 0.053328 \end{bmatrix}$$

We normalize the first part and take the scalar product.

$$\gamma_3 = \frac{1}{0.152544} \cdot \begin{bmatrix} 0.089628 \\ 0.053328 \end{bmatrix} = \begin{bmatrix} 0.67400881 \\ 0.32599119 \end{bmatrix}$$

2) *Calculating ξ* : We calculate ξ by

$$\xi_t = ((\mathbb{1}^T \alpha_T)^{-1} \cdot A) \circ (\alpha_t \otimes (\beta_{t+1} \circ B_{y_{t+1}})^T)$$

We start by calculating $((\mathbb{1}^T \alpha_T)^{-1} \cdot A)$: From before, we have

$$(\mathbb{1}^T \alpha_T)^{-1} = \frac{1}{0.152544} = 6.996$$

We have:

$$A = \begin{bmatrix} 0.6 & 0.4 \\ 0.5 & 0.5 \end{bmatrix}$$

We get:

$$8.996 \cdot A = \begin{bmatrix} 6.996 \cdot 0.6 & 6.996 \cdot 0.4 \\ 6.996 \cdot 0.5 & 6.996 \cdot 0.5 \end{bmatrix} = \begin{bmatrix} 4.198 & 2.798 \\ 3.498 & 3.498 \end{bmatrix}$$

We can now calculate $\alpha_1 \otimes (\beta_2 \circ B_{y2})^T$. We have :

$$\alpha_1 = \begin{bmatrix} 0.56 \\ 0.08 \end{bmatrix}, \quad \beta_2 = \begin{bmatrix} 0.58 \\ 0.55 \end{bmatrix}, \quad B_{y2} = \begin{bmatrix} 0.3 \\ 0.6 \end{bmatrix}$$

We calculate $\beta_2 \circ B_{y2}$:

$$\beta_2 \circ B_{y2} = \begin{bmatrix} 0.58 \\ 0.55 \end{bmatrix} \circ \begin{bmatrix} 0.3 \\ 0.6 \end{bmatrix} = \begin{bmatrix} 0.174 \\ 0.33 \end{bmatrix}$$

Outer product:

$$\begin{aligned} \alpha_1 \otimes (\beta_2 \circ B_{y2})^T &= \begin{bmatrix} 0.56 \\ 0.08 \end{bmatrix} \otimes [0.174 \quad 0.33] \\ &= \begin{bmatrix} 0.09744 & 0.1848 \\ 0.01392 & 0.0264 \end{bmatrix} \end{aligned}$$

We can now calculate ξ_1

$$\xi_1 = \begin{bmatrix} 4.198 & 2.798 \\ 3.498 & 3.498 \end{bmatrix} \circ \begin{bmatrix} 0.09744 & 0.1848 \\ 0.01392 & 0.0264 \end{bmatrix}$$

$$\xi_1 = \begin{bmatrix} 0.38325991 & 0.03650094 \\ 0.60572687 & 0.08653241 \end{bmatrix}$$

At t=2:

We have:

$$B_{y1} = \begin{bmatrix} 0.7 \\ 0.4 \end{bmatrix}, \quad \alpha_2 = \begin{bmatrix} 0.1074 \\ 0.1584 \end{bmatrix}, \quad \beta_3 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

Hadamard product for $\beta_3 \circ B_{y1}$

$$\beta_3 \circ B_{y1} = \begin{bmatrix} 1 \\ 1 \end{bmatrix} \circ \begin{bmatrix} 0.7 \\ 0.4 \end{bmatrix} = \begin{bmatrix} 0.7 \\ 0.4 \end{bmatrix}$$

Outer product:

$$\alpha_2 \otimes [0.7 \quad 0.4] = \begin{bmatrix} 0.07518 & 0.04296 \\ 0.11088 & 0.06336 \end{bmatrix}$$

We can now calculate ξ_2 :

$$\xi_2 = \begin{bmatrix} 4.198 & 2.798 \\ 3.498 & 3.498 \end{bmatrix} \circ \begin{bmatrix} 0.07518 & 0.04296 \\ 0.11088 & 0.06336 \end{bmatrix}$$

$$\xi_2 = \begin{bmatrix} 0.07341938 & 0.06873304 \\ 0.03726872 & 0.0523348 \end{bmatrix}$$

At t=3:

We have:

$$B_{y1} = \begin{bmatrix} 0.7 \\ 0.4 \end{bmatrix}, \quad \alpha_3 = \begin{bmatrix} 0.089628 \\ 0.053328 \end{bmatrix}, \quad \beta_3 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

Hadamard product for $\beta_3 \circ B_{y1}$

$$\beta_3 \circ B_{y1} = \begin{bmatrix} 1 \\ 1 \end{bmatrix} \circ \begin{bmatrix} 0.7 \\ 0.4 \end{bmatrix} = \begin{bmatrix} 0.7 \\ 0.4 \end{bmatrix}$$

Outer product:

$$\alpha_3 \otimes [0.7 \quad 0.4] = \begin{bmatrix} 0.062740 & 0.035852 \\ 0.037329 & 0.021331 \end{bmatrix}$$

We can now calculate ξ_3 :

$$\xi_3 = \begin{bmatrix} 4.198 & 2.798 \\ 3.498 & 3.498 \end{bmatrix} \circ \begin{bmatrix} 0.062740 & 0.035852 \\ 0.037329 & 0.021331 \end{bmatrix}$$

$$\xi_3 = \begin{bmatrix} 0.2839837 & 0.09127753 \\ 0.13480176 & 0.06519824 \end{bmatrix}$$

D. Update values

$$\hat{\pi} = \gamma_1 = \begin{bmatrix} 0.86784141 \\ 0.1321589 \end{bmatrix}$$

$$\hat{A} = (\mathbb{1} \otimes \gamma) \bullet \xi$$

$$\hat{B} = (\mathbb{1} \otimes \gamma) \bullet \left(\sum_{t=1}^T \gamma_t \otimes \mathbb{1}_{y_t}^T \right)$$

When referring to γ , we use the sum of the probabilities:

$$\gamma = \sum_{t=1}^T \gamma_t$$

and ξ :

$$\xi = \sum_{t=1}^T \xi_t$$

We therefore calculate:

$$\gamma = \begin{bmatrix} 0.86784141 \\ 0.1321589 \end{bmatrix} + \begin{bmatrix} 0.42888609 \\ 0.57111391 \end{bmatrix} + \begin{bmatrix} 0.67400881 \\ 0.32599119 \end{bmatrix} = \begin{bmatrix} 1.97073631 \\ 1.02926369 \end{bmatrix}$$

And

$$\xi = \begin{bmatrix} 0.38325991 & 0.03650094 \\ 0.60572687 & 0.08653241 \end{bmatrix} + \begin{bmatrix} 0.07341938 & 0.06873304 \\ 0.03726872 & 0.0523348 \end{bmatrix}$$

$$+ \begin{bmatrix} 0.2830837 & 0.09127753 \\ 0.13480176 & 0.06519824 \end{bmatrix} = \begin{bmatrix} 0.739763 & 0.19651152 \\ 0.77779736 & 0.20406545 \end{bmatrix}$$

We can now calculate

$$\mathbb{1} \otimes \gamma = \begin{bmatrix} 1 \\ 2.0923 \\ 1 \\ 1.1352 \end{bmatrix}$$

We can now calculate \hat{A}

$$\hat{A} = \begin{bmatrix} 1 \\ 2.0923 \\ 1 \\ 1.1352 \end{bmatrix} \bullet \begin{bmatrix} 0.9897 & 0.7370 \\ 0.5670 & 0.3888 \end{bmatrix} = \begin{bmatrix} 0.37537391 & 0.19092437 \\ 0.39467348 & 0.198226353 \end{bmatrix}$$

We calculate \hat{B} We first calculate the sum of the outer products:

$$\sum_{t=1}^T \gamma_t \otimes \mathbb{1}_{y_t}^T$$

At t = 1:

$$\gamma_1 \otimes [1 \quad 0] = \begin{bmatrix} 0.86784141 \\ 0.1321589 \end{bmatrix} \otimes [1 \quad 0] = \begin{bmatrix} 0.86784141 & 0.13215859 \\ 0 & 0 \end{bmatrix}$$

At t = 2:

$$\gamma_2 \otimes [0 \quad 1] = \begin{bmatrix} 0.42888609 \\ 0.57111391 \end{bmatrix} \otimes [0 \quad 1] = \begin{bmatrix} 0 & 0 \\ 0.42888609 & 0.57111391 \end{bmatrix}$$

At t = 3:

$$\gamma_3 \otimes [1 \quad 0] = \begin{bmatrix} 0.67400881 \\ 0.32599119 \end{bmatrix} \otimes [1 \quad 0] = \begin{bmatrix} 0.67400881 & 0.32599119 \\ 0 & 0 \end{bmatrix}$$




Fig. 1. B-matrix representation in ADD




Fig. 3. A-matrix representation in ADD caption reduced




Fig. 2. A-matrix representation in ADD




Fig. 4. π -matrix representation in ADD

We summarize these to get:

$$\begin{bmatrix} 0.86784141 & 0.13215859 \\ 0 & 0 \end{bmatrix} + \begin{bmatrix} 0 & 0 \\ 0.42888609 & 0.57111391 \end{bmatrix} +$$

$$\begin{bmatrix} 0.67400881 & 0.32599119 \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} 1.54185022 & 0.45814978 \\ 0.42888609 & 0.57111391 \end{bmatrix}$$

$$\hat{b} = \begin{bmatrix} 1 \\ 2.0923 \\ 1 \\ 1.1352 \end{bmatrix} \cdot \begin{bmatrix} 1.54185022 & 0.45814978 \\ 0.42888609 & 0.57111391 \end{bmatrix}$$

$$= \begin{bmatrix} 0.78237266 & 0.23247645 \\ 0.41669214 & 0.55487618 \end{bmatrix}$$

E. ADD representation

As we only need one bit to represent the the rows and columns with one bit, we only need one variable for the them, as x_1 is the variable for rows and y_1 is the variable for column.

We first make the matrices into ADD representation.

We can now use the ADD representation to calculate α and β .

When using ADD's it is important to remember, if we need to take a row from a matrix, we fix the input to the ADD by setting the x-variables to the desired row. An example is taking the third row of a matrix with 8 rows, we set, $x_1 = 1, x_2 = 1, x_3 = 0$ and $x_4 = 0$. if we need to take the second column, we set $y_1 = 1, y_2 = 0$ and $y_3 = 0, y_4 = 0$. Hadamard product is row-wise multiplication of the matrices. So to calculate the Hadamard product of two matrices, we set the x-variables to the same row in both matrices and multiply the corresponding nodes in the ADDs. To calculate a Hadamard product in ADD,

we multiply the corresponding nodes in the ADDs, as shown in the following figure.

Matrix multiplication is done by fixing the input to the first matrix and the output to the second matrix. We then sum the result of the Hadamard product of the rows of the first matrix and the columns of the second matrix. This is shown in the following figure.

ACRONYMS

AAU Aalborg University. 1

APPENDIX A

COMPILING IN DRAFT

You can also compile the document in draft mode. This shows todos, and increases the space between lines to make space for your supervisors feedback.




Fig. 5. Matrix A in ADD




Fig. 6. Matrix B in ADD




Fig. 7. Hadamard product of A and B in ADD




Fig. 8. Matrix multiplication of A and B in ADD




Fig. 9. Scalar product in ADD




Fig. 10. Kroneker product in ADD




Fig. 11. Hadamard division of A and B in ADD




Fig. 12. Katri-Rao in ADD




Fig. 13. transpose in ADD